arXiv:0704.2147v1 [nucl-th] 17 Apr 2007

A SIMULTANEOUS CENTER–OF–MASS CORRECTION OF NUCLEON DENSITY AND MOMENTUM DISTRIBUTIONS IN NUCLEI

(accepted to Ukr.Phys.J)

A. SHEBEKO, P. GRYGOROV[1]

NSC "Kharkiv Institute of Physics and Technology"
(1, Akademicheskaya Str., Kharkiv 61108, Ukraine; e-mail: shebeko@kipt.kharkov.ua),

[1]**Department of Physics and Technology, V.N. Karazin Kharkiv National University**
(31, Kurchatova Ave., Kharkiv 61108, Ukraine; e-mail: pavel_grigorov@mail.ru)

The approach exposed in the recent paper (A. Shebeko, P. Papakonstantinou, E. Mavrommatis, Eur. Phys. J. A **27**, 143 (2006)) has been applied in studying center-of-mass motion effects on the nucleon density and momentum distributions in nuclei. We use and develop the formalism based upon the Cartesian or boson representation, in which the coordinate and momentum operators are expressed through the creation and annihilation operators for oscillator quanta in the three different space directions. We are focused upon effects due to the center-of-mass and short-range nucleon correlations embedded in translationally invariant ground-state wavefunctions. The latter are constructed in the so-called fixed center-of-mass approximation, starting with a Slater determinant wave function modified by some correlator (*e.g.*, after Jastrow or Villars). It is shown how one can simplify evaluation of the corresponding expectation values that determine the distributions. The analytic expressions derived here involve the own "Tassie-Barker" factors for each distribution. As an illustration, numerical calculations have been carried out for the nucleus ^4He with the Slater determinant to describe the nucleon $(1s)^4$ configuration composed of single-particle orbitals which differ from harmonic oscillator ones at small distances. Such orbitals simulate somewhat short-range repulsion between nucleons. Special attention is paid to a simultaneous shrinking of the center–of–mass corrected density and momentum distributions compared to the purely $(1s)^4$ shell nontranslationally invariant ones.

1. Introduction

Treatment of the center–of–mass (CM) motion has been an attractive subject of exploration in earlier and more recent studies of nuclear theory (see, *e.g.*, [1]– [9]). Those studies originated from the necessity to remedy a deficiency of the nuclear many-body wave function (WF), namely its lack of translational invariance (TI) wherever shell-model single-particle (s.p.) WF's are used for its construction. This deficiency is important in quite a number of cases.

In the present investigation we adopt the "fixed-CM approximation" [10, 11] as a recipe to restore TI of a many-body WF which does not possess this property. We apply it when evaluating the elastic form factor (FF) $F(q)$ and the nucleon momentum distribution (MD) $\eta(p)$ for light nuclei, and more specifically for ^4He in its ground state (g.s.). Following [9] we prefer to deal with the intrinsic quantities which are determined as expectation values of appropriate (multiplicative) operators that depend on the corresponding Jacobi variables and act on the intrinsic WF's. We have seen in [9] that the intrinsic density distribution (DD) $\rho_{int}(r)$, being defined by the Fourier transform of $F(q)$, does not coincide with the diagonal part of the one-body density matrix (1DM), which is related in a standard manner to the intrinsic MD. In the context, we also note that the term "one-body" used here is somewhat conventional. Let us mention that $F(q)$ and $\eta(p)$ can be related to the different quantities measured via electron–nucleus collisions, respectively, the elastic electron scattering cross sections and the inclusive electron scattering cross sections. First of all, we mean comparatively simple relations in the Born approximation with the plane electron waves. In addition, to the so–called approximation of small interaction times (see [12]– [14] and refs. therein) the double differential (e, e') reaction cross section becomes proportional to an integral of $\eta(p)$ over the momentum range that is fixed with certain combination (the so-called y - scaling variable) of the momentum transfer q and the energy transfer ω (cf. [15]). Of course, in the framework of these approximations one neglects off–shell effects in the electron scattering on bound nucleons and meson exchange currents (MEC) contributions to an effective electromagnetic (e.m.) interaction with nuclei. The latter should be taken into account (see, *e.g.*, refs. [16,17]) when describing the electron scattering on nuclei, especially at high momentum transfers (in particular, helping to remove certain discrepancy between theory and experiment in the vicinity of the first minimum $|F_{ch}(q)|$ at $q^2 = 10 fm^{-2}$ for ^4He). Therefore, any comparison with experimental data omitting such physical inputs has a restricted character. Nevertheless, in case of light nuclei every approximate evaluation of intrinsic quantities, being independent of different constraints originated from reaction mechanisms, can be compared with microscopic ("exact") results. In the respect, our addressing to the alpha particle seems to be perfectly explicable.

The aim of this paper is to show to what extent the approach developed in refs. [9,12,18,19] can be useful in calculations with more realistic WF's than simple harmonic oscillator model (HOM) ones. In this connection, we consider the CM correction of $F(q)$ and $\eta(p)$ treated on an equal physical footing, *viz.*, by using one and the same translationally invariant g.s. WF that incorporates the nucleon-nucleon short-range correlations (SRC). One should note that despite much interest over the last two decades concerning the MD in nuclei [20]– [26], its CM correction does not appear to have been properly treated except in certain studies, where harmonic oscillator (HO) wave functions were used (see, *e.g.*, [12]). Note also calculations beyond HOM in [27]. The underlying formalism with basic definitions is exposed in the following section. Section 3 contains analytic results of our derivations beyond HOM, while the corresponding numerical results are discussed and compared with experimental data in section 4.

2. The intrinsic form factor, density and momentum distributions with short-range correlations included

By definition, the intrinsic (elastic) FF of a nonrelativistic system with the mass number A and the total angular momentum equal to zero is

$$F(q) = F_{int}(q) \equiv \langle \Phi_{int} \mid \exp[\imath \vec{q} \cdot (\hat{\vec{r}}_1 - \hat{\vec{R}})] \mid \Phi_{int} \rangle, \qquad (1)$$

where Φ_{int} is the intrinsic WF of the system (nucleus), $\hat{\vec{r}}_1$ the coordinate operator for nucleon number 1, and $\hat{\vec{R}} = A^{-1} \sum_{i=1}^{A} \hat{\vec{r}}_i$ the CM operator.

In the fixed-CM approximation, according to the Ernst, Shakin and Thaler (EST) prescription [11] the nuclear many-body WF with the total momentum $\vec{P}$ can be written in the form:

$$\mid \Psi_P \rangle = |\vec{P}) \mid \Phi_{int}^{EST} \rangle, \qquad (2)$$

where a round bracket is used to represent a vector in the space of the CM coordinate,so that $|\vec{P})$ means the eigenstate of total momentum operator $\hat{\vec{P}}$. The intrinsic WF after EST

$$\mid \Phi_{int}^{EST} \rangle = \frac{(\vec{R} = 0 \mid \Phi\rangle}{[\langle \Phi \mid \vec{R} = 0)(\vec{R} = 0 \mid \Phi\rangle]^{1/2}} \qquad (3)$$

is constructed from an arbitrary (in general, translationally non-invariant) WF Φ, by requiring that the CM coordinate $\vec{R}$ be equal to zero. The corresponding FF is the ratio

$$F_{EST}(q) = \frac{A(q)}{A(0)},$$

$$A(q) = \langle \Phi \mid (2\pi)^3 \delta(\hat{\vec{R}}) \exp[i\vec{q} \cdot (\hat{\vec{r}}_1 - \hat{\vec{R}})] \mid \Phi \rangle. \qquad (4)$$

Using the Cartesian representation in which

$$\hat{\vec{r}} = \frac{r_0}{\sqrt{2}}\,(\hat{\vec{a}}^\dagger + \hat{\vec{a}}),\ \hat{\vec{p}} = \imath\frac{p_0}{\sqrt{2}}\,(\hat{\vec{a}}^\dagger - \hat{\vec{a}}) \tag{5}$$

with the Bose commutation rules,

$$[\hat{a}_i, \hat{a}_j] = 0, \qquad [\hat{a}_i, \hat{a}_j^\dagger] = \delta_{ij} \quad (i,j = 1,2,3) \tag{6}$$

and arbitrary real c-numbers r_0 and p_0 that meet the condition

$$r_0 p_0 = 1, \tag{7}$$

one can show (see [9], [19] and Appendix A) that

$$A(q) = \exp\left(-\frac{\bar{r}_0^2 q^2}{4}\right) U(q), \tag{8}$$

$$U(q) = \int d\vec{\lambda} \exp\left(-\frac{r_0^2 \lambda^2}{4A}\right) F(\vec{v}, \vec{s}), \tag{9}$$

$$F(\vec{v}, \vec{s}) = \langle \Phi | \hat{O}_1(\vec{v} + \vec{s}\ )\hat{O}_2(\vec{v}) \ldots \hat{O}_A(\vec{v}) | \Phi \rangle, \tag{10}$$

where

$$\hat{O}_\gamma(\vec{x}) = \exp(-\vec{x}^* \hat{\vec{a}}_\gamma^\dagger) \exp(\vec{x}\hat{\vec{a}}_\gamma) \equiv \hat{E}_\gamma^\dagger(-\vec{x})\hat{E}_\gamma(\vec{x}) \tag{11}$$

$(\gamma = 1, \ldots, A)$

with

$$\vec{s} = \imath\frac{r_0}{\sqrt{2}}\ \vec{q},\ \ \vec{v} = \imath\frac{r_0}{\sqrt{2}A}(\vec{\lambda} - \vec{q}) \tag{12}$$

and the renormalized "length"parameter

$$\bar{r}_0 = \sqrt{\frac{A-1}{A}} r_0.$$

Further, starting from the definition of the intrinsic MD (see [9]),

$$\eta(p) = \langle \Phi_{int} \mid \delta(\hat{\vec{p}}_1 - \hat{\vec{P}}/A - \vec{p}) \mid \Phi_{int} \rangle, \tag{13}$$

we consider the distribution in the fixed-CM approximation:

$$\eta_{EST}(p) = \frac{\langle \Phi \mid (2\pi)^3 \delta(\hat{\vec{R}}) \delta(\hat{\vec{p}}_1 - \hat{\vec{P}}/A - \vec{p}) \mid \Phi \rangle}{\langle \Phi \mid (2\pi)^3 \delta(\hat{\vec{R}}) \mid \Phi \rangle} \tag{14}$$

and the Fourier transform

$$\eta_{EST}(p) = (2\pi)^{-3} \int \exp(-\imath\vec{p}\vec{x}) N(x)/N(0) d\vec{x} \tag{15}$$

with

$$N(x) = \langle \Phi \mid (2\pi)^3 \delta(\vec{R}) \exp[\imath(\vec{p}_1 - \vec{P}/A)\vec{x}] \mid \Phi \rangle. \tag{16}$$

We see the certain resemblance between the structure functions $N(x)$ and $A(q)$, *viz.*, both are determined by the expectation values of similar multiplicative operators with one and the same trial WF Φ. Owing to this, using the same algebraic technique we get

$$N(x) = \exp\left(-\frac{\bar{p}_0^2 x^2}{4}\right) D(x), \tag{17}$$

$$D(x) = \int d\vec{\lambda} \exp\left(-\frac{r_0^2 \lambda^2}{4A}\right) F(\vec{v}\,', \vec{s}\,'), \tag{18}$$

where

$$\vec{s}\,' = -\frac{p_0}{\sqrt{2}}\vec{x}, \quad \vec{v}\,' = \frac{\imath r_0}{\sqrt{2}A}(\vec{\lambda} - \imath p_0^2 \vec{x}) \tag{19}$$

and

$$\bar{p}_0 = \sqrt{\frac{A-1}{A}} p_0.$$

Certain relation of the MD to the corresponding intrinsic density matrix has been shown in [9].

After this let us assume a trial WF,

$$\mid \Phi \rangle = \mid \Phi_{corr} \rangle = \hat{C}(1, 2, \cdots, A) \mid Det \rangle \tag{20}$$

with the Slater determinant

$$\mid Det \rangle = \frac{1}{\sqrt{A!}} \sum_{\hat{\mathcal{P}} \in S_A} \epsilon_{\mathcal{P}} \hat{\mathcal{P}} \{ \mid \phi_{p_1}(1) \rangle \cdots \mid \phi_{p_A}(A) \rangle \}. \tag{21}$$

Here $\epsilon_{\mathcal{P}}$ is the parity factor for the permutation $\mathcal{P}$, ϕ_a the occupied orbital with the quantum numbers $\{a\}$ and the summation runs over all permutations of the symmetric group S_A.

The A-particle operator $\hat{C} = \hat{C}(\hat{\vec{r}}_\alpha - \hat{\vec{r}}_\beta,\ \hat{\vec{p}}_\alpha - \hat{\vec{p}}_\beta)$ [1] introduces the SRC and meets all necessary requirements of the translational and Galileo invariance, the permutable and rotational symmetry, etc. However, being translationally invariant itself such a model introduction of correlations does not enable to restore the TI violated with such a shell-model WF as the Slater determinant.

---

[1] Of course, the operator may be spin and isospin dependent

What it follows can be used with the Jastrow correlator [28]

$$\hat{C} = \prod_{\alpha<\beta}^{A} f(\hat{\vec{r}}_{\alpha\beta}), \tag{22}$$

where $f(\hat{\vec{r}}_{\alpha\beta})$ is a two-body correlation factor whose deviation from unity occurs only for small distances $r_{\alpha\beta} = |\vec{r}_\alpha - \vec{r}_\beta|$ less than a correlation radius r_c.

Another popular option goes back to the lectures by Villars in [29] (see also [30]) with a unitary operator

$$\hat{C} = \exp(-\imath\hat{G}), \tag{23}$$

$$\hat{G} = \sum_{\alpha<\beta} \hat{g}(\alpha,\beta), \tag{24}$$

where the Hermitian operator $\hat{g}(\alpha,\beta)$ acts onto the space of the pair (α,β). In particular, we could follow the simplest Darmstadt ansatz [31]:

$$\hat{g}(\alpha,\beta) = \frac{1}{2}\{\vec{s}\ (\hat{\vec{r}}_{\alpha\beta})\hat{\vec{p}}_{\alpha\beta} + \hat{\vec{p}}_{\alpha\beta}\vec{s}\ (\hat{\vec{r}}_{\alpha\beta})\}, \tag{25}$$

where $\vec{s}$ is a function of the relative coordinate $\hat{\vec{r}}_{\alpha\beta} = \hat{\vec{r}}_\alpha - \hat{\vec{r}}_\beta$. Its canonically conjugate momentum $\hat{\vec{p}}_{\alpha\beta} = \frac{1}{2}(\hat{\vec{p}}_\alpha - \hat{\vec{p}}_\beta)$.

Keeping in mind similar constructions we rewrite expectation (10) as

$$F(\vec{v},\vec{s}) = \langle\Phi(-\vec{v}) \mid \hat{E}_1^\dagger(-\vec{s})\hat{E}_1(\vec{s}) \mid \Phi(\vec{v})\rangle, \tag{26}$$

where

$$\mid \Phi(\vec{x})\rangle = \hat{E}_1(\vec{x})\dots\hat{E}_A(\vec{x}) \mid \Phi\rangle,$$

since $\hat{E}_1(\vec{v}+\vec{s}) = \hat{E}_1(\vec{v})\hat{E}_1(\vec{s})$ and $[\hat{E}_\alpha(\vec{x}),\hat{E}_\beta(\vec{y})] = 0$ $(\alpha,\beta = 1,\dots,A)$ for any vectors $\vec{x}$ and $\vec{y}$.

Moreover, we find that

$$\hat{E}(\vec{x})\ \hat{\vec{r}}\ \hat{E}^{-1}(\vec{x}) = \hat{\vec{r}} + \frac{r_0}{\sqrt{2}}\ \vec{x} \tag{27}$$

and

$$\hat{E}(\vec{x})\hat{\vec{p}}\ \hat{E}^{-1}(\vec{x}) = \hat{\vec{p}} - \imath\frac{p_0}{\sqrt{2}}\ \vec{x}. \tag{28}$$

Remind that $E^\dagger \neq E^{-1}$. In other words, $\hat{E}_\alpha(\vec{x})$ is the displacement operator in the space of nucleon states with the label α.

Due to this property when handling the similarity transformation

$$\hat{C}' = \hat{E}_1(\vec{x})\dots\hat{E}_A(\vec{x})\hat{C}(\hat{\vec{r}}_\alpha - \hat{\vec{r}}_\beta,\ \hat{\vec{p}}_\alpha - \hat{\vec{p}}_\beta)\times$$

$$\times\hat{E}_1^{-1}(\vec{x})\dots\hat{E}_A^{-1}(\vec{x}),$$

we get

$$\hat{C}' = \hat{C}(\hat{E}_\alpha(\vec{x})\hat{\vec{r}}_\alpha\hat{E}_\alpha^{-1}(\vec{x}) - \hat{E}_\beta(\vec{x})\hat{\vec{r}}_\beta\hat{E}_\beta^{-1}(\vec{x}),$$

$$\hat{E}_\alpha(\vec{x})\hat{\vec{p}}_\alpha\hat{E}_\alpha^{-1}(\vec{x}) - \hat{E}_\beta(\vec{x})\hat{\vec{p}}_\beta\hat{E}_\beta^{-1}(\vec{x})) =$$

$$= \hat{C}(\vec{r}_\alpha - \vec{r}_\beta, \vec{p}_\alpha - \vec{p}_\beta) = \hat{C}$$

i.e.,

$$\hat{C}' = \hat{C}. \tag{29}$$

Recall that C is a function of all the relative coordinates and their canonically conjugate momenta.
From eqs. (20) and (29) it follows that

$$\mid \Phi_{corr}(\vec{x})\rangle \equiv \hat{E}_1(\vec{x})\dots\hat{E}_A(\vec{x}) \mid \Phi_{corr}\rangle =$$

$$= \hat{C} \mid Det(\vec{x})\rangle. \tag{30}$$

Here $\mid Det(\vec{x})\rangle = \hat{E}_1(\vec{x})\dots\hat{E}_A(\vec{x}) \mid Det\rangle$ is a new Slater determinant composed of the renormalized orbitals,

$$\mid \phi_a(\vec{x};\alpha)\rangle = \hat{E}_\alpha(\vec{x}) \mid \phi_a(\alpha)\rangle \quad (\alpha = 1,\dots,A), \tag{31}$$

viz.,

$$\mid Det(\vec{x})\rangle = \frac{1}{\sqrt{A!}}\sum_{\hat{\mathcal{P}}\in S_A} \epsilon_{\mathcal{P}}\hat{\mathcal{P}}\{\mid \phi_{p_1}(\vec{x};1)\rangle \cdots \mid \phi_{p_A}(\vec{x};A)\rangle\}. \tag{32}$$

In their turn, such orbitals can be evaluated in a concise analytic form as initial ones are linear combinations of the HOM orbitals (see Appendix A).

Following (26) we arrive to

$$F_{corr}(\vec{v},\vec{s}) \equiv \langle \Phi_{corr}(-\vec{v}) \mid \hat{E}_1^\dagger(-\vec{s})\hat{E}_1(\vec{s}) \mid \Phi_{corr}(\vec{v})\rangle =$$

$$= \langle Det(-\vec{v}) \mid \hat{C}^\dagger \hat{E}_1^\dagger(-\vec{s})\hat{E}_1(\vec{s})\hat{C} \mid Det(\vec{v})\rangle. \qquad (33)$$

Expressions (8) and (17) with expectations $F(\vec{v},\vec{s})$ and $F(\vec{v}\,',\vec{s}\,')$, which are determined by eq. (33), are certain base for our calculations.

2.1 Several working formulae: application to ^4He

In special case of the pure HOM $(1s)^4$ configuration occupied by the four nucleons in ^4He we have

$$\mid \Phi_{corr}(\vec{x})\rangle = \mid \Phi_{corr}\rangle = \hat{C} \mid (1s)^4\rangle, \qquad (34)$$

taking into account that the HOM g.s. $\mid (1s)^4\rangle$ is the vacuum for operators $\hat{\vec{a}}_\alpha$ ($\alpha = 1,\ldots,A$). It is the case, where $\mid Det(\vec{v})\rangle$ does not depend on $\vec{v}$ coinciding with the initial Slater determinant $\mid (1s)^4\rangle$. Hence,

$$F_{1s}(\vec{v},\vec{s}) = \langle (1s)^4 \mid \hat{C}^\dagger \hat{E}_1^\dagger(-\vec{s})\hat{E}_1(\vec{s})\hat{C} \mid (1s)^4\rangle. \qquad (35)$$

In other words, under such a simplification the function $F(\vec{v},\vec{s})$ in integral (9) becomes independent of $\vec{\lambda}$ and we get

$$U(q) = U_{1s}(q)\int \exp\left(-\frac{r_0^2\lambda^2}{4A}\right) d\vec{\lambda},$$

so that

$$\frac{U_{1s}(q)}{U_{1s}(0)} = \frac{\langle (1s)^4 \mid \hat{C}^\dagger \hat{E}_1^\dagger(-\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{E}_1(\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{C} \mid (1s)^4\rangle}{\langle (1s)^4 \mid \hat{C}^\dagger\hat{C} \mid (1s)^4\rangle}. \qquad (36)$$

Thus, the FF of interest is

$$F_{EST}(q) = F_{TB}(q)F_{IPM}(q)F_{corr}(q), \qquad (37)$$

where according to eq.(A.1) we have the Tassie-Barker $F_{TB}(q)$ and the HOM FF $F_{HOM}(q)$. The factor

$$F_{corr}(q) = \frac{\langle (1s)^4 \mid \hat{C}^\dagger \hat{E}_1^\dagger(-\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{E}_1(\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{C} \mid (1s)^4\rangle}{\langle (1s)^4 \mid \hat{C}^\dagger\hat{C} \mid (1s)^4\rangle} \qquad (38)$$

incorporates the SRC in any way.

At this point, one can proceed, at least, along the two guidelines. One of them could be based upon the representation

$$\langle (1s)^4 \mid \hat{C}^\dagger \hat{E}_1^\dagger(-\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{E}_1(\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{C} \mid (1s)^4\rangle =$$

$$= \langle (1s)^4 \mid \hat{C}_1^\dagger(-\vec{q})\hat{C}_1(\vec{q}) \mid (1s)^4\rangle, \qquad (39)$$

where

$$\hat{C}_1(\vec{q}) = \hat{E}_1(\imath\frac{r_0}{\sqrt{2}}\vec{q})\hat{C}(\hat{\vec{r}}_1,\hat{\vec{p}}_1,\cdots)\hat{E}_1^{-1}(\imath\frac{r_0}{\sqrt{2}}\vec{q}) =$$

$$= C\left(\hat{\vec{r}}_1 + \imath\frac{\vec{q}}{2}r_0^2, \hat{\vec{p}}_1 + \frac{\vec{q}}{2}, \cdots\right). \qquad (40)$$

Other continuation is prompted by the relation

$$\hat{E}_1^\dagger(-\imath\frac{r_0}{\sqrt{2}}\;\vec{q}\;)\hat{E}_1(\imath\frac{r_0}{\sqrt{2}}\;\vec{q}\;) = \exp\left(\frac{r_0^2q^2}{4}\right)\exp(\imath\vec{q}\;\hat{\vec{r}}_1),$$

that gives rise to

$$F_{corr}(q) = \exp\left(\frac{r_0^2q^2}{4}\right)F_C(q), \qquad (41)$$

$$F_C(q) = \frac{\langle (1s)^4 \mid \hat{C}^\dagger \exp(\imath\vec{q}\hat{\vec{r}}_1)\hat{C} \mid (1s)^4\rangle}{\langle (1s)^4 \mid \hat{C}^\dagger\hat{C} \mid (1s)^4\rangle}, \qquad (42)$$

where $F_C(q)$ is the no CM corrected FF with the correlated g.s. $\hat{C} \mid (1s)^4\rangle$.

Analogously, we find

$$N(x) = N_{TB}(x)\,N_{HOM}(x)\,N_{corr}(x) \qquad (43)$$

with the own Tassie-Barker factor

$$N_{TB}(x) = \exp\left(\frac{p_0^2\;x^2}{4A}\right) \qquad (44)$$

and

$$N_{HOM}(x) = \exp\left(-\frac{p_0^2\;x^2}{4}\right), \qquad (45)$$

$$N_{corr}(x) = \frac{\langle (1s)^4 \mid \hat{C}^\dagger \hat{E}_1^\dagger(\frac{p_0}{\sqrt{2}}\vec{x})\hat{E}_1(-\frac{p_0}{\sqrt{2}}\vec{x})\hat{C} \mid (1s)^4\rangle}{\langle (1s)^4 \mid \hat{C}^\dagger\hat{C} \mid (1s)^4\rangle}. \tag{46}$$

Again, different continuations are possible (cf. the transition from eq. (38) to eqs. (39) and (41)). In particular, with the help of

$$\hat{E}_1^\dagger(\frac{p_0}{\sqrt{2}}\vec{x})\hat{E}_1(-\frac{p_0}{\sqrt{2}}\vec{x}) = \exp\left(\frac{p_0^2\, x^2}{4}\right)\exp\left(\imath\hat{\vec{p}}_1\vec{x}\right)$$

we get

$$N_{corr}(x) = \exp\left(\frac{p_0^2\, x^2}{4}\right) N_C(x), \tag{47}$$

$$N_C(x) = \frac{\langle (1s)^4 \mid \hat{C}^\dagger \exp(\imath\hat{\vec{p}}_1\vec{x})\hat{C} \mid (1s)^4\rangle}{\langle (1s)^4 \mid \hat{C}^\dagger\hat{C} \mid (1s)^4\rangle}. \tag{48}$$

The Fourier transform

$$\eta_C(p) = \frac{1}{(2\pi)^3}\int \mathrm{e}^{-\imath\vec{p}\vec{x}} N_C(x)d\vec{x} \tag{49}$$

gives us the one-body momentum distribution (OBMD) without the CM correction of the model g.s. $\hat{C} \mid (1s)^4\rangle$.

By definition, the intrinsic DD is

$$\rho_{int}(r) = \frac{1}{(2\pi)^3}\int \mathrm{e}^{-\imath\vec{q}\vec{r}} F_{int}(q)d\vec{q} =$$

$$= \langle \Phi_{int} \mid \delta(\hat{\vec{r}}_1 - \hat{\vec{R}} - \vec{r}) \mid \Phi_{int}\rangle, \tag{50}$$

so that the relations

$$\rho_{EST}(r) = \frac{1}{(2\pi)^3}\int \mathrm{e}^{-\imath\vec{q}\vec{r}} F_{EST}(q)d\vec{q}$$

and

$$\rho_C(r) = \frac{1}{(2\pi)^3}\int \mathrm{e}^{-\imath\vec{q}\vec{r}} F_C(q)d\vec{q}$$

are, respectively, the one-body density distribution (OBDD) with the CM correction and the no CM corrected distribution.

Thus, we have shown (with the help of purely algebraic means) that evaluation of the distributions can be reduced to the well-known treatment. Indeed, expectations values (42) and (48) occur in all conventional calculations with the many-particle WF (20), *i.e.,* without any CM correction. Diverse methods have been elaborated when evaluating similar quantities (see, *e.g.,* [6], [8], [30], [31], [32], [33], [34], [35] and refs. therein). In this work we confine ourselves to comparatively simple computations for a $(1s)^4$ configuration, where a short-range repulsion between nucleons is introduced in an effective way, *viz.,* modifying the s.p. orbital as in [10]. Respectively, the following WF is used in the next section.

3. Analytic expressions for the form factor, density and momentum distributions with the single-particle wave function beyond HOM

In accordance with [10] we employ the normalized Radhakant, Khadkikar and Banerjee (RKB) radial orbital for the lowest s.p. state of ^4He,

$$\phi^{RKB}(r) = \frac{1}{\sqrt{1+\beta^2}}(\phi_{00}(r) + \beta\phi_{10}(r)), \tag{51}$$

where ϕ_{00} and ϕ_{10} are the normalized HO radial eigenfunctions:

$$\phi_{00}(r) = 2\sqrt{\frac{1}{\sqrt{\pi}b_H}}\frac{r}{b_H}\exp\left(-\frac{r^2}{2b_H^2}\right), \tag{52}$$

$$\phi_{10}(r) = \sqrt{\frac{3!}{\sqrt{\pi}b_H}}\frac{r}{b_H}\left[1 - \frac{2}{3}\frac{r^2}{b_H^2}\right]\exp\left(-\frac{r^2}{2b_H^2}\right) \tag{53}$$

for the states with $n = 0, l = 0$ and $n = 1, l = 0$, respectively. Here b_H is the HO parameter and β is a mixing parameter.

The RKB WF allows one to obtain the following expressions for the density distribution (normalized to unity), for the point proton FF as well as for the MD (also normalized to unity):

$$\rho_{sp}^{RKB}(r) = \frac{1}{(\sqrt{\pi}b_H)^3(1+\beta^2)}\exp\left(-\frac{r^2}{b_H^2}\right)\times$$

$$\times \left[1+\sqrt{\frac{3}{2}}\beta\left(1-\frac{2r^2}{3b_H^2}\right)\right]^2, \tag{54}$$

$$F_{sp}^{RKB}(q)=\frac{1}{1+\beta^2}\exp\left(-\frac{(b_Hq)^2}{4}\right)\times$$

$$\times\left[1+\beta^2+\frac{\beta}{\sqrt{6}}\left(1-\sqrt{\frac{2}{3}}\beta\right)b_H^2q^2+\frac{\beta^2b_H^4q^4}{24}\right], \tag{55}$$

$$\eta_{sp}^{RKB}(p)=\frac{b_H^3}{\pi\sqrt{\pi}(1+\beta^2)}\exp(-b_H^2p^2)\times$$

$$\times \left[1-\sqrt{\frac{3}{2}}\beta\left(1-\frac{2}{3}b_H^2p^2\right)\right]^2. \tag{56}$$

3.1 The CM corrected form factor $F(q)$ *and its reduction to quadratures*

Assuming a Slater determinant as the g.s. $(1s)^4$ of ^4He its FF in the fixed-CM approximation can be written in the form (cf. [10]):

$$F_{EST}(q)=\frac{\int F_{1s}(|\vec{q}+\vec{u}|)F_{1s}^3(u)d\vec{u}}{\int F_{1s}^4(u)d\vec{u}}, \tag{57}$$

where

$$F_{1s}(v)\equiv\int \mathrm{e}^{i\vec{v}\vec{r}}\phi_{1s}^2(r)\mathrm{d}\vec{r}=\frac{4\pi}{v}\int\phi_{1s}^2(r)\sin vr\ r\mathrm{d}r$$

is the no CM-corrected FF.

In case of the RKB–like s.p. WFs whose orbitals are truncated expansions in the radial HO eigenfunctions the multiple integrals in the r.h.s. of eq.(63) can be expressed through simple integrals. The respective algebraic technique has been developed in ref. [12] and exposed recently in [37] (see also Appendix A to the present paper). Its application with the RKB orbital enables us to get

$$F^{RKB}(q)=\frac{A^{RKB}(q)}{A^{RKB}(0)}, \tag{58}$$

where

$$A^{RKB}(q)=I_1(q)+I_2(q),$$

$$I_1(q)=\frac{4\pi}{qb_H^4}\exp\left(-\frac{3}{16}q^2b_H^2\right)\times$$

$$\times\int_0^\infty\{B_2\left[\frac{1}{4}(t-\frac{3}{4}\ qb_H)^2\right]M_2^3\left[\frac{1}{4}(t+\frac{1}{4}\ qb_H)^2\right]-$$

$$-B_2\left[\frac{1}{4}(t+\frac{3}{4}\ qb_H)^2\right]M_2^3\left[\frac{1}{4}(t-\frac{1}{4}\ qb_H)^2\right]\}\exp(-t^2)tdt,$$

$$I_2(q)=\frac{\pi}{b_H^3}\exp\left(-\frac{3}{16}q^2b_H^2\right)\times$$

$$\times\int_0^\infty\{B_2\left[\frac{1}{4}(t-\frac{3}{4}qb_H)^2\right]M_2^3\left[\frac{1}{4}(t+\frac{1}{4}qb_H)^2\right]+$$

$$+B_2\left[\frac{1}{4}(t+\frac{3}{4}qb_H)^2\right]M_2^3\left[\frac{1}{4}(t-\frac{1}{4}qb_H)^2\right]\}\exp(-t^2)dt,$$

$$A^{RKB}(0)=\frac{4\pi}{b_H^3}\int_0^\infty M_2^4\left[\frac{1}{4}\ t^2\right]\exp(-t^2)t^2dt.$$

The functions $M_2(z)$ and $B_2(z)$ are second degree polynomials of the variable z



Fig.1. The point-like FF (left) and the charge FF (right) of ^4He. Curves calculated with RKB WF using the EST prescription (solid) and without the CM-correction (dashed); experimental points from [39]. Other clarifications in the text

$$M_2(z)=m_0+m_1z+m_2z^2$$

$$B_2(z)=h_0+h_1z+h_2z^2,$$

where the constants are related to the mixing parameter β

$$m_0=1+\beta^2,$$

$$m_1=2\sqrt{2/3}\beta(1-\sqrt{2/3}\beta),$$

$m_2 = (2/3)\beta^2,$

and

$h_0 = m_0 + m_1 + 2m_2,$

$h_1 = m_1 + 2m_2,$

$h_2 = m_2.$



Fig.2. The one-body density distribution (OBDD) for $(1s)^4$ configuration with RKB orbital in the fixed-CM approximation (solid) and without the CM correction (dashed). For two sets of parameters: $b_H = 0.8532$ fm and $\beta = -0.4738$ (top); $b_H = 0.8532$ fm and $\beta = 0$ (bottom). Dot-dashed with the parametrization from [41]

3.2 The CM corrected momentum distribution $\eta(p)$ and its reduction to quadratures

In parallel, starting from eq.(16), we obtain for the $(1s)^4$ configuration with the Slater determinant $\mid \Phi\rangle = \mid (1s)^4\rangle$ (see Appendix B to Lect.I in [37]):

$$N_{EST}(x) = \int \mathrm{d}\vec{k}\, \langle 1s \mid \exp\left(\imath \frac{\vec{k}\hat{\vec{r}}}{A}\right) \exp\left(\imath \frac{A-1}{A}\hat{\vec{p}}\vec{x}\right) \mid 1s\rangle \times$$

$$\times \langle 1s \mid \exp\left(\imath \frac{\vec{k}\hat{\vec{r}}}{A}\right) \exp\left(-\imath \frac{\hat{\vec{p}}\vec{x}}{A}\right) \mid 1s\rangle^3. \qquad (59)$$



Fig.3. The one-body momentum distribution (OBMD) for the $(1s)^4$ configuration with RKB orbital. Difference between the curves is the same as in fig. 2. Dot-dashed with parametrization from [42]

Again, using the representation (5) and splitting the exponents involved in eq.(67) with the succesive normal ordering of the operators $\hat{\vec{a}}^{\,\dagger}$ and $\hat{\vec{a}}$ (the former are to the left from the latter), one can show that

$$\frac{N(x)^{RKB}}{N(0)^{RKB}} = \exp\left(-\frac{A-1}{A}\frac{x^2}{4b_H^2}\right)\frac{J(x)}{J(0)}, \qquad (60)$$

where the integral $J(x)$ is determined by

$$J(x) = \int\limits_0^\infty \exp\left(-\frac{r_0^2\lambda^2}{4A}\right) g(\lambda^2; x^2)\lambda^2 d\lambda,$$

$$g(\lambda^2; x^2) = \left(A_1 + \frac{1}{3}A_2\right)B_1^3 + \left(A_1 + \frac{3}{5}A_2\right)B_1^2 B_2 +$$

$$+3\left(\frac{1}{5}A_1 + \frac{1}{7}A_2\right)B_1 B_2^2 + \left(\frac{1}{7}A_1 + \frac{1}{9}A_2\right)B_2^3,$$

$$A_1 = 1 + \beta^2 - \sqrt{\frac{2}{3}}\,\beta\left[1 + \sqrt{\frac{2}{3}}\beta\right]\left(\frac{A-1}{A}\right)^2 \frac{p_0^2 x^2}{2} +$$

$$+\sqrt{\frac{2}{3}}\beta\left[1 - \sqrt{\frac{2}{3}}\beta\right]\frac{r_0^2\lambda^2}{2A^2} +$$

$$+\frac{1}{6}\beta^2\left[\left(\frac{A-1}{A}\right)^2 \frac{p_0^2 x^2}{2} - \frac{r_0^2\lambda^2}{2A^2}\right]^2,$$

$$A_2 = \frac{1}{6}\beta^2 \left(\frac{A-1}{A}\right)^2 \frac{x^2\lambda^2}{A^2},$$

$$B_1 = 1 + \beta^2 - \sqrt{\frac{2}{3}}\beta \left[1 + \sqrt{\frac{2}{3}}\beta\right] \frac{p_0^2 x^2}{2A^2} +$$

$$+\sqrt{\frac{2}{3}}\beta \left[1 - \sqrt{\frac{2}{3}}\beta\right] \frac{r_0^2\lambda^2}{2A^2} + \frac{1}{6}\beta^2 \left[\frac{p_0^2 x^2}{2A^2} - \frac{r_0^2\lambda^2}{2A^2}\right]^2,$$

$$B_2 = \frac{1}{6}\beta^2 \frac{x^2\lambda^2}{A^4}$$

Thus, the structure function $N^{RKB}(x)$ can be reduced to one-dimensional integrals similar to those derived for $F^{RKB}(q)$. Here $A = 4$, but we allow A to be changeable, particularly, in order to check that the corresponding distribution

$$\eta_{EST}^{RKB}(p) = \frac{1}{2\pi^2 p} \int\limits_0^\infty N^{RKB}(x)/N^{RKB}(0)\sin(px)x dx$$

to the limit $A \to \infty$ yields the no CM corrected distribution (62).

4. Results and discussion

Analytic expressions obtained in sect.2 for the density and momentum distributions and their Fourier transforms are sufficiently general to be applied in different translationally invariant treatments with the SRC included. The corresponding formulae derived in sect.3 in case of the ^4He nucleus have been employed to carry out our calculations beyond the simple HOM. Their numerical results are displayed in figs.1-4.



Fig.4. Variations $log\ \eta_{EST}^{RKB}(p)$ (solid), $log\ \eta_{sp}^{RKB}(p)$ (dashed) and $log\ \eta_{Morita}(p)$ (dot-dashed) with p. Notation $log\ \eta_{Morita}(p) = \left(\frac{4}{3}\right)^3 W^{SN}(\frac{4}{3}p)$, where function $W^{SN}(x)$ calculated with a convenient parametrization from [42]. Points are resulted from [24]

In fig. 1 we show our calculation of the charge FF, $F_{ch}(q^2) = f_p(q)F(q)$ of the alpha particle, using eq.(55) and eq.(58) and considering for the finite proton size factor $f_p(q)$ the Chandra and Sauer prescription [40]. The two parameters b_H and β have been determined by the least-square fitting to the experimental values [39]: their best-fit values are $b_H = 0.8532$ fm and $\beta = -0.4738$ ($\chi^2 \simeq 13.07$). These values have been utilized in our calculations shown in figs. 2-4.

As it is seen in fig.1 (its left part), the CM correction leads to a considerable qualitative change of q-dependence of the FF: its first minimum and second maximum are shifted towards higher q-values. This difference between solid and dashed curves is due to the different behavior of the respective densities at small distances $r \leq 1$ fm: see fig. 2, top, where the dashed curve is $\rho_{sp}^{RKB}(r)$ by eq.(54), while the solid is determined by

$$\rho^{RKB}(r) = \frac{1}{2\pi^2 r} \int\limits_0^\infty F^{RKB}(q)\sin(qr)q dq. \tag{61}$$

Moreover, we see that each of the distributions (for the simple HOM orbital on the bottom and for the RKB orbital on the top), after being CM corrected, increases in its central but decreases in its peripheral region. One may say that we encounter a specific effect of shrinking the OBDD owing to the translationally invariant treatment.

In addition, there is a central depression of the density distribution (cf. the upper and lower dashed lines in fig.2). Such a change is not unexpected since the RKB WF represents a simple way to allow for some of the effects of short-range repulsion between the nucleons in ^4He. These numerical results get an explicit confirmation if one writes

$$\rho_{sp}^{RKB}(0) = \frac{1}{(\sqrt{\pi}b_H)^3}\frac{1}{1+\beta^2}\left(1+\sqrt{\frac{3}{2}}\beta\right)^2 . \qquad (62)$$

Evidently, the inequality

$$\rho_{sp}^{RKB}(0) \le \rho_{sp}^{RKB}(0)\mid_{\beta=0} \equiv \rho^{HOM}(0) = \frac{1}{(\sqrt{\pi}b_H)^3}$$

takes place for negative β values with $\mid \beta \mid < 2\sqrt{6}$.

In parallel, we show in fig. 3 that the corresponding change of the OBMD has much in common with that for the OBDD, *viz.*, the distribution $\eta_{EST}^{RKB}(p)$ turns out to be shrunk in the above sense relative to the distribution $\eta_{sp}^{RKB}(p)$. Thus, we see a simultaneous shrinking of the density distribution $\rho(r)$ and the momentum distribution $\eta(p)$. As has been shown in [12] (see also [14]), such a simultaneous change of these distributions plays a substantial role in getting a fair treatment of the data on the elastic and inelastic electron scattering of ^4He. Let us recall that there the charge FF and the dynamic FF of ^4He were calculated using one and the same HOM WF, corrected both with the fixed-CM approximation and the Peierls-Yoccoz prescription [44].

Regarding the properties of these simultaneously corrected distributions in detail, we would like to emphasize a practical consequence of their interpretation. This aspect becomes especially transparent in the case of the simple HOM $(1s)^4$ configuration, where we have

$$\rho_{EST}^{HOM}(r) = [\sqrt{\pi}\bar{r}_0]^{-3}\exp(-r^2/\bar{r}_0^2)$$

vs

$$\rho_{sp}^{HOM}(r) = [\sqrt{\pi}r_0]^{-3}\exp(-r^2/r_0^2)$$

and

$$\eta_{EST}^{HOM}(p) = [\sqrt{\pi}\bar{p}_0]^{-3}\exp(-p^2/\bar{p}_0^2)$$

vs

$$\eta_{sp}^{HOM}(p) = [\sqrt{\pi}p_0]^{-3}\exp(-p^2/p_0^2)$$

Thus, the inclusion of CM corrections gives rise to the two independent renormalizations, $r_0 \equiv b_H \rightarrow \bar{r}_0 = \sqrt{3/4}r_0$ and $p_0 \equiv {b_H}^{-1} \rightarrow \bar{p}_0 = \sqrt{3/4}p_0$, of the oscillatory parameter values, r_0 and p_0 (cf. [12]). Evidently, such changes are not equivalent to a hasty replacement of p_0 by $\sqrt{4/3}p_0$ if one follows the Tassie-Barker recipe with $b_H \rightarrow \sqrt{3/4}b_H$ only.

Now, following the conventional way of determining the HOM parameter r_0, as in ref. [12], we will use the expansions

$$F_{ch}(q^2) = 1 - \frac{1}{6}q^2r_{ch}^2 + \cdots ,$$

$$f_p(q) = 1 - \frac{1}{6}q^2r_p^2 + \cdots$$

and

$$F(q) = 1 - \frac{1}{6}q^2r_{rms}^2 + \cdots ,$$

where we have in HOM

$$r_{rms}^2 = \frac{3}{2}r_0^2,$$

so that

$$r_{ch}^2 = \frac{3}{2}r_0^2 + r_p^2,$$

whence

$$r_0^2 \equiv r_{exp}^2 = \frac{2}{3}\left[r_{ch}^2 - r_p^2\right] . \qquad (63)$$

Doing so for the CM corrected quantities we find the similar relation

$$\bar{r}_0^2 = \frac{2}{3}\left[r_{ch}^2 - r_p^2\right] = r_{exp}^2 \qquad (64)$$

with the identical q-dependence $F_{EST}^{HOM}(q) = F^{HOM}(q) = \exp(-q^2 r_{exp}^2/4)$. At the same time the difference between the respective OBMDs becomes more considerable than after the substitution $r_0 \to \sqrt{\frac{A}{A-1}}r_{exp}$ in $\eta_{sp}^{HOM}(p) = \frac{r_0^3}{\pi^{3/2}}\exp(-p^2r_0^2)$ that gives

$$\eta_{sp}^{HOM}(p) = \left(\frac{A}{A-1}\right)^{3/2}\frac{r_{exp}^3}{\pi^{3/2}}\times$$

$$\times\exp\left[-\frac{A}{A-1}p^2r_{exp}^2\right]$$

vs

$$\eta_{sp}^{HOM}(p) = \frac{r_{exp}^3}{\pi^{3/2}}\exp(-p^2r_{exp}^2).$$

Under the simultaneous CM correction of the OBDD and OBMD we have

$$\eta_{EST}^{HOM}(p) = \left(\frac{A}{A-1}\right)^{3}\frac{r_{exp}^3}{\pi^{3/2}}\times$$

$$\times\exp\left[-\left(\frac{A}{A-1}\right)^2p^2r_{exp}^2\right] \tag{65}$$

vs

$$\eta_{sp}^{HOM}(p) = \frac{r_{exp}^3}{\pi^{3/2}}\exp(-p^2r_{exp}^2), \tag{66}$$

that is equivalent to the substitution $r_0 \to \frac{A}{A-1}r_{exp}$ in $\eta_{sp}^{HOM}(p) = \frac{r_0^3}{\pi^{3/2}}\exp(-p^2r_0^2)$.

Note also that the product $\bar{r}_0\bar{p}_0 = 1-A^{-1} \neq 1$, unlike the relation $r_0p_0 = 1$. In this connection, following [19] let us remind the commutation rules for intrinsic coordinate $\vec{r}\,' = \vec{r}-\vec{R}$ and conjugate momenta $\vec{p}\,' = \vec{p}-\vec{P}/A$

$$[\vec{r}\,'_l, \vec{p}\,'_j] = i\delta_{l,j}(1-1/A), \quad (l,j=1,2,3) \tag{67}$$

One can show that the corresponding uncertainty principle is related to the deviation from unity. Thus, the uncertainty principle does not contradict the simultaneous shrinking of the density and momentum distributions (see also [37], Lect.I, Suppl. C)

In case of the RKB function we get

$$r_{rms}^2 = \frac{3}{2}r_0^2 - \frac{\beta\sqrt{6}}{1+\beta^2}\left(1-\sqrt{\frac{2}{3}}\beta\right)r_0^2. \tag{68}$$

It means that the short-range repulsion involved in the WF with a negative β leads to some increasing the rms radius, *viz.*, $r_{rms}^{RKB} > r_{rms}^{HOM}$. For the values $b_H = 0.8532$ fm and $\beta = -0.4738$ the formula (68) yields $r_{rms}^{RKB} = 1.429$ fm, so that the corresponding charge radius is equal to $r_{ch}^{RKB} = 1.667$ fm. Here we employ the charge proton radius $r_p = 0.86$ fm (see, for example, Appendix 7 in [43]).

The CM correction gives an opposite effect. Indeed, after some calculation we find that $r_{rms}^{EST} = 1.309$ fm for the same $b_H = 0.8532$ fm and $\beta = -0.4738$. From this it follows that $r_{ch}^{EST} = 1.566$ fm.

The variation of $\log\eta_{EST}^{RKB}$ and of $\log\eta_{sp}^{RKB}$ with p is depicted in fig.4 for a wider range of momenta. It is seen from fig.4 that the allowance of the CM motion improves the description of the available data on the OBMD of the alpha particle. It is further seen from fig.4 that in the translationally invariant quantity according to the fixed-CM prescription the "seagull" behavior appearing in the variation of the corresponding s.p. one becomes somewhat less pronounced. The dip is diminished and it moves to smaller values of momentum.

Finally, we would like to point out in connection to the comparison with the s.p. distributions that the CM corrected OBDD and OBMD become closer to the corresponding microscopic ones by using their convenient parametrizations from [41] and [42], as one can see in figs.3 and 4. According to the communication [42] one has to introduce the factor $(2\pi)^{-3}$ to reproduce the momentum distribution $\eta_{Morita}(p)$ which is one of the significant results obtained by the Sapporo group. At this point, let us recall that these authors employed the so-called ATMS-method, where ATMS is abbreviation "Amalgamation of Two-body correlations into Multiple Scattering process", to construct the variational WF of the ^4He nucleus (see [21] and refs. therein). Along the variational approach a considerable progress was made when including more dynamics of the realistic nucleon-nucleon interaction such as the effect of its tensor component (cf. [22]).

5. Concluding remarks

We have seen how the approach exposed in [9] can be extended to the translationally invariant evaluation of the density and momentum distributions in nuclei. The present analysis shows that the restoration of translational invariance in the Slater determinant WF of ^4He by means of the fixed-CM correction (the EST prescription) gives rise as a whole to essential changes in the r-, p- and q-dependences of the OBDD $\rho(r)$, the OBMD $\eta(p)$ and the charge FF $F_{ch}(q)$, respectively. We have seen that the correlation between nucleons induced by the fixation of the center-of-mass of the nucleus results in the simultaneous shrinking of $\rho(r)$ and $\eta(p)$. Meanwhile, this effect has been revealed here beyond the pure HOM extending the available experience.

Also, this study demonstrates the relative importance of the CM and SRC corrections for the same nucleus, *viz.*, the shrinking of the density and momentum distributions owing to the use of translationally invariant g.s. wave functions of ^4He vs their broadening after the inclusion of short-range repulsion in these wave functions at small distances $r < 1$ fm. It is true that the latter has been introduced in our calculations in a simple manner. Nevertheless, there are all reasons to believe that the algebraic method employed here might be helpful within more sophisticated approaches, where the short-range correlations are taking into account via the Jastrow factor or other correlation operator (see, *e.g.* [31]). At present, the corresponding applications are in progress both for the ^4He and ^{16}O nuclei.

Acknowledgements

One of us, A.S., is very grateful to Michael Grypeos and his colleagues for their hospitality during his visits to Department of Theoretical Physics at Aristotle University of Thessaloniki, where this work was begun.

Also, it is a great pleasure for us to thank A. Antonov and M. Gaidarov for sending the numerical values of the momentum distribution displayed by the points in fig.4.

Appendix A

Some details of calculations beyond HOM

Here we want to illustrate a convenient method for evaluation of the expectations in question being aimed at some general (model-independent) results (cf. [9], [19]).

First of all, we have by recurring the Cartesian representation:

$$
\begin{aligned}
&\exp\left[\imath\vec{q}\,(\hat{\vec{r}}_1-\hat{\vec{R}})\right]= \\
&=\exp\left[\imath\vec{q}\,\left(\frac{A-1}{A}\right)\,\hat{\vec{r}}_1\right]\exp\left[-\imath\vec{q}\,\frac{\hat{\vec{r}}_2}{A}\right]\cdots= \\
&=F_{TB}(q)\;F_{HOM}(q)\times \\
&\times\exp\left[\imath\vec{q}\left(\frac{A-1}{A}\right)\frac{r_0}{\sqrt{2}}\;\hat{\vec{a}}_1^{\dagger}\right]\times \\
&\times\exp\left[\imath\vec{q}\left(\frac{A-1}{A}\right)\frac{r_0}{\sqrt{2}}\;\hat{\vec{a}}_1\right]\times \\
&\qquad\times\exp\left[-\imath\vec{q}\frac{r_0}{\sqrt{2}A}\;\hat{\vec{a}}_2^{\dagger}\right]\exp\left[-\imath\vec{q}\frac{r_0}{\sqrt{2}A}\;\hat{\vec{a}}_2\right]\ldots, \qquad (A.1)
\end{aligned}
$$

$F_{TB}(q)=\exp(\frac{r_0^2q^2}{4A})$, $F_{HOM}(q)=\exp(-\frac{r_0^2q^2}{4})$, where the index α at $\hat{\vec{a}}_\alpha(\hat{\vec{a}}_\alpha^{\dagger})$ is the individual particle number $(\alpha=1,\cdots,A)$.

Thereat, the Tassie-Barker factor $F_{TB}(q)$ appears automatically due to a specific structure of the operators involved. In other words, its appearance is independent of any nuclear properties (in general, properties of a finite system). The only mathematical tool that has been used is the Baker-Hausdorff relation:

$$
\mathrm{e}^{\mathrm{A+B}}=\mathrm{e}^{\mathrm{A}}\;\mathrm{e}^{\mathrm{B}}\;\mathrm{e}^{-\frac{1}{2}[\mathrm{A,B}]}, \qquad (A.2)
$$

that is valid with arbitrary operators $\hat{A}$ and $\hat{B}$ for which the commutator $\left[\hat{A},\hat{B}\right]$ commutes with each of them. Further, applying eq.(A.2) in combination with

$$
(2\pi)^3\,\delta\left(\hat{\vec{R}}\right)=\int\exp\left(\imath\vec{\lambda}\hat{\vec{R}}\right)d\vec{\lambda}, \qquad (A.3)
$$

one can show that the expectation value $A(q)$ in eq. (4) and the expectation value $N(x)$ in eq. (16) are expressed through one and the same function $F(\vec{x}, \vec{y})$ that depends, respectively, on the arguments $\vec{x} = \vec{v}$, $\vec{y} = \vec{s}$ (as in eq.(9)) and $\vec{x} = \vec{v'}$, $\vec{y} = \vec{s'}$ (as in eq.(18)). In other words, we have constructed the common generating function for each of them. One should stress that this result has been obtained independently of the model WF Φ.

The algebraic technique shown here turns out to be useful for practical calculations with the Slater determinants like $\mid \Phi\rangle$ (see [18]) or the Slater determinants modified by different correlators (for instance, the Jastrow factor).

In the simplest case of the independent particle model (IPM) $(1s)^4$ configuration for ^4He with the Slater determinant $\mid \Phi\rangle = \mid (1s)^4\rangle$ we get omitting the nonessential factor $[A!]^{-1}$,

$$A^{IPM}(q) = \exp\left[-\frac{\bar{r}_0^2 q^2}{4}\right] U^{IPM}(q), \qquad (A.4)$$

$$U^{IPM}(q) = \int d\vec{\lambda} \exp\left[-\frac{r_0^2 \lambda^2}{4A}\right] f(\vec{\lambda}, \vec{q}\ ), \qquad (A.5)$$

$$f(\vec{\lambda}, \vec{q}) = \langle 1s \mid \exp(-\vec{\alpha}^* \hat{\vec{a}}^\dagger) \exp(\vec{\alpha}\ \hat{\vec{a}}) \mid 1s\rangle \times$$
$$\times \langle 1s \mid exp(-\vec{\beta}^* \hat{\vec{a}}^\dagger) \exp(\vec{\beta}\vec{a}) \mid 1s\rangle^3, \qquad (A.6)$$

$$\vec{\alpha} = \imath \frac{r_0}{\sqrt{2}A}[\vec{\lambda} + (A-1)\vec{q}],\ \vec{\beta} = \imath \frac{r_0}{\sqrt{2}A}[\vec{\lambda} - \vec{q}],$$

with the renormalized "length"parameter

$$\bar{r}_0 = \sqrt{\frac{A-1}{A}}\ r_0$$

In our case $r_0 = b_H$. We are keeping the notations with a mass number A $(= 4)$ to point out certain trend in the A-dependence.

At this point, let us note that the RKB orbital (or another model orbital) being composed of the basis states of the spherical representation can be written then as a superposition of the basis states $\mid n_1 n_2 n_3\rangle$ of the Cartesian representation (see, *e.g.*, the monograph [38] and refs. therein),

$$\mid n_1 n_2 n_3\rangle = \frac{1}{\sqrt{n_1! n_2! n_3!}} (\hat{\vec{a}}_1^{\ \dagger})^{n_1} (\hat{\vec{a}}_2^{\ \dagger})^{n_2} (\hat{\vec{a}}_3^{\ \dagger})^{n_3} \mid 0\rangle, \qquad (A.7)$$

where the vector $\mid 0\rangle \equiv \mid 000\rangle$ is the vacuum state with respect to the destruction operators $\hat{a}_i$ $(i = 1, 2, 3)$, *e.g.*,

$$\hat{\vec{a}} \mid 0\rangle = 0. \qquad (A.8)$$

It is proved that for the RKB-orbital,

$$\mid 1s\rangle = [1 + \beta^2]^{-1/2} [1 - (\beta/\sqrt{6})\ \hat{\vec{a}}^{\ \dagger} \hat{\vec{a}}^{\ \dagger}\ ] \mid 0\rangle. \qquad (A.9)$$

Substituting (A.9) into (A.6) (when calculating the ratio $A^{IPM}(q)/A^{IPM}(0)$, the normalization factor $[1+\beta^2]^{-1/2}$ can be omitted) we find

$$\exp(\vec{\chi} \cdot \vec{a}) \mid 1s\rangle = [1 - (\beta/\sqrt{6})(\hat{\vec{a}}^{\ \dagger} + \vec{\chi})(\hat{\vec{a}}^{\ \dagger} + \vec{\chi})] \mid 0\rangle \qquad (A.10)$$

for any complex vector $\vec{\chi}$.

Now, after modest effort we obtain

$$\langle 1s \mid \exp(-\vec{\chi}^{\ *} \cdot \hat{\vec{a}}^{\ \dagger}) \exp(\vec{\chi} \cdot \hat{\vec{a}}) \mid 1s\rangle =$$
$$= 1 + \beta^2 - \frac{2}{3}\ \beta^2 \vec{\chi}^{\ *} \vec{\chi} -$$
$$- \frac{\beta}{\sqrt{6}}\ [\vec{\chi}^{\ *} \vec{\chi}^{\ *} + \vec{\chi}\ \vec{\chi}\ ] + \frac{\beta^2}{6}\ (\vec{\chi}^{\ *} \vec{\chi}^{\ *})(\vec{\chi}\ \vec{\chi}) \qquad (A.11)$$

From (A.11) it follows, for instance,

$$\langle 1s \mid \exp(\imath \vec{\alpha}^* \hat{\vec{a}}^{\ \dagger}) \exp(\imath \vec{\alpha} \hat{\vec{a}}) \mid 1s\rangle = M_2 \left(\frac{\vec{\alpha}^* \vec{\alpha}}{2}\right), \quad (A.12)$$

where the polynomial $M_2(z)$ is given at the end of subsect. 3.1.

1. J.L. Friar, Nucl. Phys. A **173**, 257 (1971).
2. C. Ciofi degli Atti, L. Lanto and P. Toropainen, Phys. Lett. B **42**, 443 (1972).
3. C.M. Vincent, Phys. Rev. C **8**, 929 (1973).
4. A.E.L. Dieperink, T. de Forest,Jr., Phys. Rev. C **10**, 543 (1974).
5. D. Van Neck *et al.*, Phys. Rev. C **57**, 2308 (1998).
6. Bogdan Mihaila, Jochen Heisenberg, Phys. Rev. C **60**, 054303 (1999).
7. K.W. Schmid, Eur. Phys. J. A **12**, 29 (2001); ibid., **13**, 319 (2002); ibid., **14** 413 (2002); ibid., **16**, 475 (2003).
8. P. Navratil, Phys. Rev. C **70**, 014317 (2004).
9. A. Shebeko, P. Papakonstantinou, E. Mavrommatis, Eur. Phys. J. A **27**, 143 (2006).

10. S. Radhakant, S.B. Khadkikar, B. Banerjee, Nucl. Phys. A **142**, 81 (1970).
11. D. Ernst, C. Shakin, R. Thaler, Phys. Rev. C **7**, 925 (1973); ibid., 1340.
12. S. Dementiji, V. Ogurtzov, A. Shebeko, Sov. J. Nucl. Phys. **22**, 6 (1976).
13. A.Yu. Korchin, A.V. Shebeko, Z. Phys. A **321**, 687 (1985).
14. S. V. Dementiji *et al.*, J. Phys. Soc. Jap. **57**, 2988 (1988).
15. C. Ciofi degli Atti, E. Pace, G. Salme In: *Proceedings of the 2nd Workshop on Perspectives in Nuclear Physics at Intermediate Energies* (WS, Singapore, 1985).
16. T. Katayama, Y. Akaishi, H. Tanaka, Prog. Theor. Phys. **67** 236 (1982).
17. R. Schiavilla, V.R. Pandharipande, R.B. Wiringa, Phys. Rev. C **41**, 309 (1990).
18. A. Shebeko, N. Goncharov, Sov. J. Nucl. Phys. **18**, 532 (1974).
19. A. Korchin, A. Shebeko, Ukr. J. Phys. **22** (10), 1646 (1977); arXiv: nucl-th/0601014.
20. S. Frullani, J. Mougey, Adv. Nucl. Phys., eds. J. Negele and E. Vogt, Vol. **14** (New York, 1984).
21. Y.Akaishi, Inter. Rev. Nucl. Phys. Vol. **4** (WS, Singapore, 1986).
22. R. Schiavilla, V.R. Pandharipande, R.B. Wiringa, Nucl. Phys. A **449**, 219 (1986).
23. A.N. Antonov, P.E. Hodgson, J.Zh. Petkov, *Nucleon Momentum and Density Distributions in Nuclei* (Clarendon Press, Oxford, 1988).
24. C. Ciofi degli Atti, E. Pace, G. Salme, Phys. Rev. C **43**, 1155 (1991).
25. K. Ypsilantis, M. Grypeos, J. Phys. G.:Nucl. Part. Phys. **15**, 397 (1989).
26. K. Ypsilantis, M. Grypeos, J. Phys. G.: Nucl. Part. Phys. **21**, 1701 (1995).
27. M. Grypeos, C. Koutroulos, A. Shebeko, K. Ypsilantis, Part. Nucl. Lett. **2 [111]**, 74 (2002).
28. R. Jastrow, Phys. Rev. **98**, 1497 (1955).
29. F. M. H. Villars in: *Rendiconti S.I.F.*, XXIII Corso (New York, 1963).
30. J. Da Providencia, C. M. Shakin, Ann. Phys. **30**, 95 (1964).
31. H. Feldmeier *et al.*, Nucl. Phys. A **632**, 61 (1998).
32. M. Gaudin, *et al.*, Nucl. Phys. A **176**, 237 (1971).
33. M. Dal Ri, S. Stringari, O. Bohigas, Nucl. Phys. A **376**, 81 (1982).
34. Ch. C. Moustakidis *et al.*, Phys. Rev C **64**, 014314 (2001).
35. M. Alvioli, C. Ciofi degli Atti, H. Morita , Phys. Rev. C **72**, (2005); arXiv: nucl-th/054310.
36. H. Bateman, A. Erdélyi, *Tables of Integral Transforms.* V.**1** (McGraw-Hill, 1954).
37. A. Shebeko, *Lectures on Selected Topics of Nuclear Theory* (Thessaloniki, 2000).
38. V. Neudachin, Yu. Smirnov, *Nucleon Clusters in Light Nuclei* ("Nauka", Moscow, 1964).
39. R.F. Frosch *et al.*, Phys. Rev. **160**, 1308 (1966);
    R.G. Arnold *et al.*, Phys. Rev. Lett. **40**, 1429 (1978).
40. H. Chandra, G. Sauer, Phys. Rev. C **13**, 245 (1976).
41. H. Morita, Y. Akaishi, O. Endo, H. Tanaka, Prog. Theor. Phys. **78**, 1117 (1987).
42. H. Morita, Y. Akaishi, H. Tanaka, Prog. Theor. Phys. **79**, 863 (1988); H. Morita, Private communication to A. Shebeko.
43. T. Ericson, W. Weise, *Pions and Nuclei*, (Clarendon Press, Oxford, 1988).
44. R. Peierls, J. Yoccoz, Proc. Phys. Soc. A **70**, 381 (1957).